

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 03-01-2003		2. REPORT TYPE Technical Paper		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE New Application of Bromotrimethylsilane: Elaboration of Aldehydes/Ketones into Homologous α,β -Unsaturated Esters via β -Hydroxy Esters				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Suri Suresh and Jacob C. Marcischak				5d. PROJECT NUMBER 4847	
				5e. TASK NUMBER 0244	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRSA 10 E. Saturn Blvd. Edwards AFB, CA 93524-7680				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-PR-ED-TP-2002-305	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-PR-ED-TP-2002-305	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
20030213 111					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT A	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Leilani Richardson
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code) (661) 275-5015

✓DTS

MEMORANDUM FOR PRS (In-House/Contractor Publication)

FROM: PROI (STINFO)

16 Dec 2002

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2002-305**
Suri, Suresh (PRSA) and Marcischak, Jacob C. (ERC) "New Application of Bromotrimethylsilane:
5952 Elaboration of Aldehydes/Ketones into Homologated α,β -Unsaturated Esters via β -Hydroxy Esters"

Tetrahedron Letters
(Journal Article) (Deadline: N/A)

(Statement A)

New Application of Bromotrimethylsilane: Elaboration of Aldehydes/Ketones into Homologous α,β -Unsaturated Esters via β -Hydroxy Esters

Suresh C. Suri^{*¶} and Jacob C. Marcischak[§]

[¶]Air Force Research Laboratory/PRSP; [§]ERC Inc.
10 East Saturn Blvd.; Edwards Air Force Base, CA 93524

Abstract: α,β -unsaturated ester is formed when β -hydroxy-ester is reacted with bromotrimethylsilane generated from chlorotrimethylsilane-lithium bromide in acetonitrile.

The α,β -unsaturated ester is one of the important chemical functionalities that are encountered very commonly in organic synthesis. The standard method of introducing two carbon homologated α,β -unsaturated esters involves Horner-Wadsworth-Emmons¹ (HWE) modification of Wittig-olefination or Peterson-olefination² of aldehyde/ketone.

As a part of synthetic project, the conversion of β -hydroxy-ester to the corresponding β -bromo-ester was required. There is plethora of reagents³ to transform directly/indirectly alcohol into the corresponding bromo- derivative. Bromotrimethylsilane⁴ (TMSBr) was made the reagent of choice for converting hydroxy- ester to bromo- ester because of its weak electrophilicity and its inertness towards dealkylative transformation of esters, ethers, or carbamates under mild condition. Although many methods have been reported for the generation of bromotrimethylsilane, we generated bromotrimethylsilane *in situ* using chlorotrimethylsilane-lithium bromide⁵ (TMSCl-LiBr) in dry acetonitrile. In an effort to convert β -hydroxy-ester to β -bromo-ester using TMSCl-LiBr, we have observed the formation of α,β -unsaturated ester. To the best of our knowledge this type of chemical transformation by TMSBr has not been reported earlier. Herein, we report two-carbon homologation of aldehyde/ketone to α,β -unsaturated esters via β -hydroxy-ester.

To determine the generality of bromotrimethylsilane assisted chemical transformation of β -hydroxy-ester to α,β -unsaturated ester, a number of two-carbon homologated β -hydroxy-ester **8-13** were prepared from corresponding aldehyde/ketone **1-7** (TABLE-1) using lithio-




TABLE-1

Aldehyde/ketone	β -Hydroxy Ester (% Yield)	α,β -Unsaturated Ester (% Yield) ^{a,b}
Ph-CHO 1	Ph-CH(OH)CH ₂ CO ₂ Et 8 ⁶ (82)	 15 ⁷ (76) ^a
 2	 9 (92)	 16 ⁷ (73) ^a
 3	 10 ⁶ (76)	 17a (85) ^a 17b (3) ^a
 4	 11 (81)	 18 (84) ^a
 5	 12 (95)	 19 (60) ^a
 6	 13 ⁶ (50)	 20a (29) ^b 20b ⁷ (45) ^b 20c (18) ^b
 7	 14 ⁶ (70)	 21a (45) ^b 21b ⁷ (9) ^b 21c (16) ^b

a: Isolated yield; b: The yield is based on GC-Mass analysis


ethylacetate⁶ that is generated by addition of dry ethylacetate to a THF solution of lithium bis-trimethylsilylamide at $-70\text{ }^{\circ}\text{C}$ (**Scheme-1**). Thus, benzaldehyde (**1**), benzophenone (**2**), cinnamaldehyde (**3**), p-methylacetophenone (**4**), norbornan-2-one (**5**), cyclopentanone (**6**), and cycloheptanone (**7**) furnished ethyl-3-hydroxy-3-phenylpropionate (**8**)⁶, ethyl-3-hydroxy-3,3-

diphenylpropionate (**9**), ethyl-3-hydroxy-5-phenyl-pent-4-enoate (**10**)⁶, ethyl-3-hydroxy-2-*p*-tolyl butyrate (**11**), ethyl-(2-hydroxy-bicyclo[2.2.1]hept-2-yl)acetate (**12**), ethyl-(1-hydroxy-cyclopentyl)-acetate (**13**)⁶ and ethyl-(1-hydroxy-cyclohexyl)acetate (**14**)⁶ respectively in respectable yield.

When an acetonitrile solution of chlorotrimethylsilane-lithium bromide was treated with hydroxy esters **8**, **9** and **10**, only ethyl-(*E*)-cinnamate (**15**)⁷ and ethyl-5-phenyl-penta-2(*E*),4(*E*)-dienoate (**16**)⁸ and ethyl-3,3-diphenyl-2-propenoate (**17**)⁷ were formed respectively. Whereas, exposure of **11** to TMSBr furnished separable mixture of ethyl-3-*p*-tolyl-but-2(*E*)-enoate (**18**)⁹. Treatment of 2-ethylacetyl-norbornan-2-ol (**12**) with TMSBr furnished a non-separable (*E/Z*) isomer mixture of ethyl-2-norbornylidene **19**. The presence of isomeric mixture was reflected by the ¹³C NMR data¹⁰ of **19**. The hydroxy ethyl esters **13** & **14** gave mixture of unsaturated esters **20/21** (a-b) and bromo-esters **20/21**(c) respectively. It was observed by GC-mass and ¹³C NMR that the mixture **20/21** (a-c) collapsed to **20/21** (a-b) when refluxed with DBU. It was also observed that the reaction took place at room temperature in case of aromatic hydroxy esters (**8-11**) whereas it required refluxing for aliphatic hydroxy esters (**12-14**).

The chemical transformation can be rationalized by assuming silylation of β -hydroxy ester with TMSBr to furnish alkoxytrimethylsilane that in turn is hydrolyzed by *in situ* generated HBr to yield carbonium ion (Scheme-2). Elimination of proton (H^+) leads to α,β -unsaturated ester and trapping of carbonium ion by a bromo nucleophile (Br^-) furnishes corresponding bromo-compound.

Scheme-2



In conclusion, we have demonstrated a new application of bromotrimethylsilane by converting β -hydroxy ester to α,β -unsaturated esters. The above results illustrate the efficacy of TMSBr reaction with aromatic/aliphatic β -hydroxy esters having secondary and tertiary alcohol

group. This methodology could be an alternative for HWE method for the preparation of two carbon homologated α,β -unsaturated esters from aldehyde/ketone.

Acknowledgment: This financial grant from Air Force Office of Scientific Research to support this research is gratefully acknowledged.

References

1. Wadsworth, W. S. and Emmons, W. D. *J. Am. Chem. Soc.* **1961**, 83, 1733; Wadsworth, W. S. and Emmons, W. D. *Org. Syn.* **1965**, 45, 44. For comprehensive review, see: Kulkarni, Y. S. *Aldrichimica Acta* **1990**, 23, 39; Maryanoff, B. E.; Reitz, A. B. *Chem. Rev.* **1989**, 89, 863; Wadsworth, W. S., Jr. *Org. React.* 1977, 25, 73-254; Boutagy, J. and Thomas, R. *Chem. Rev.* **1974**, 74, 87; Stec, W.J. *Acc. Chem. Res.* **1983**, 16, 411-417.
2. Peterson, D. J. *J. Org. Chem.* **1968**, 33, 780; Bellassoued, M. and Ozanne, N. *J. Org. Chem.* **1995**, 60, 6582. For reviews, see: Magnus, P. *Aldrichimica Acta* **1980**, 13, 43; Agar, D. J. *Org. React.* **1990**, 38, 1-223; Agar, D. J. *Synthesis* **1984**, 384.
3. For a comprehensive list of reagents, see: Larock, R. C. *Comprehensive Organic Transformations*; VCH Publishers, Inc., New York, NY, 1989; pp 353-363.
4. Schmidt, A. H. *Aldrichimica Acta* **1981**, 14, 31; Martinelli, M. J. in *Encyclopedia of Reagents for Organic Synthesis*, ed. Paquette, L. A.; John Wiley & Sons, Inc., New York, NY; 784-786
5. Olah, G. A., Balaram Gupta, B. G., Malhotra, R., Narang, S. C. *J. Org. Chem.* **1980**, 45, 1638.
6. Rathke, M. W. *J. Am. Chem. Soc.* **1970**, 92, 3222. β -Hydroxy esters can also be prepared from α -trimethylsilylester and aldehyde/ketone in the presence of fluoride ion [Nakamura, E., Shimuza, M., and Kuwajima, I. *Tett. Lett.* **1976**, 1699]
7. Rathke, M. W. and Bouhlet, E. *Syn. Comm.* **1990**, 20, 869; Tanaka, K.; Yamagishi, N.; Tanikaga, R. and Kaji, A. *Bull. Soc. Japan* **1979**, 52, 3619.
8. Bellassoued, M. and Ozanne, N. *J. Org. Chem.* **1995**, 60, 6582.

9. Srikrishna, A. and Sundarababu, G. *Tetrahedron* **1990**, 46, 3601.
10. ^{13}C NMR (CDCl_3) data of **19**: δ 171.28, 170.14, 167.02, 109.89, 108.92, 59.58, 47.24, 42.98, 41.10, 40.01, 39.46, 39.15, 36.42, 35.59, 32.65, 31.11, 28.82, 28.58, 27.91, 27.73, 14.57.